



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Enantioselective Rhodium-Catalyzed Nucleophilic Allylation of Cyclic Imines with Allylboron Reagents

Citation for published version:

Luo, Y, Hepburn, HB, Chotsaeng, N & Lam, HW 2012, 'Enantioselective Rhodium-Catalyzed Nucleophilic Allylation of Cyclic Imines with Allylboron Reagents', *Angewandte Chemie International Edition*, vol. 51, no. 33, pp. 8309-8313. <https://doi.org/10.1002/anie.201204004>

Digital Object Identifier (DOI):

[10.1002/anie.201204004](https://doi.org/10.1002/anie.201204004)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Angewandte Chemie International Edition

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.




Enantioselective Rhodium-Catalyzed Nucleophilic Allylation of Cyclic Imines with Allylboron Reagents**

Yunfei Luo, Hamish B. Hepburn, Nawasit Chotsaeng, and Hon Wai Lam*

Enantioselective rhodium(I)-catalyzed additions of organoboron reagents to π -electrophiles are a major class of reactions for the generation of enantioenriched chiral compounds.^[1] Attractive features of these reactions include: (i) the availability of a wide range of chiral ligands that impart high enantioselectivities across several classes of electrophiles; (ii) relatively mild reaction conditions, and (iii) the availability, stability, functional group tolerance, and usually low toxicity of organoboron reagents. To date, this field has been dominated by the enantioselective addition of arylboron reagents,^[1] though there have also been reports of additions of alkenylboron^[2] and alkynylboron reagents.^[3,4,5] Notably absent are the corresponding rhodium-catalyzed enantioselective additions of allylboron reagents,^[6,7] despite the widespread importance of nucleophilic allylations in synthesis.^[8,9] In this communication, we describe the first enantioselective rhodium-catalyzed additions of allylboron reagents to π -electrophiles in the form of asymmetric allylation of cyclic imines using potassium allyltrifluoroborates. Not only do aldimines undergo highly enantioselective allylations, ketimines are also effective substrates. Furthermore, highly stereoselective additions of substituted allyltrifluoroborates are described.

The importance of chiral homoallylic amines as building blocks for chemical synthesis has led to significant efforts to develop catalytic enantioselective nucleophilic allylations of imines or their derivatives.^[8b,10,11] However, processes that employ allylboron reagents constitute a field that is still in its infancy,^[8b,10,12,13] and we therefore selected this area as a platform in which to develop a new rhodium-catalyzed variant that could offer increased substrate scope and utility. Our experiments began with attempted allylation of various benzaldehyde-derived imines using potassium allyltrifluoroborate (2 equiv)^[14,15] in the presence of 2.5 mol% of $[\text{Rh}(\text{cod})\text{Cl}]_2$ and MeOH (5 equiv) in dioxane at 80 °C (Table 1). Unfortunately, satisfactory results were not

Table 1: Rh-catalyzed allylation of benzaldehyde-derived imines.^[a]




Entry	R	1 [%] ^[b]	3 [%] ^[b]	4 [%] ^[b]
1	Ph	>95	<5	<5
2	P(O)Ph ₂	>95	<5	<5
3	SO ₂ NMe ₂	85	5	10
4	Ts	60	25	15
5	Ns	42	28	30

[a] Reactions were conducted using 0.10 mmol of **1**. [b] Determined by ¹H NMR analysis of the unpurified reaction mixtures. Ts = *p*-toluenesulfonyl, Ns = *p*-nitrobenzenesulfonyl.

obtained. With *N*-phenyl or *N*-diphenylphosphinoyl imines, only starting material was recovered (entries 1 and 2), while trace quantities of the allylation product were observed using a dimethylsulfamyl imine (entry 3). With more reactive *N*-sulfonylimines, appreciable quantities of homoallylic sulfonamides were obtained, but significant quantities of starting imine remained, along with benzaldehyde resulting from imine hydrolysis (entries 4 and 5).

In light of our recent discovery that cyclic imines are highly effective substrates for enantioselective rhodium-catalyzed alkenylations,^[16] the allylation of benzoxathiazine-2,2-dioxide **5a** was investigated (Table 2). Under conditions identical to those employed in Table 1, complete

Table 2: Rh-catalyzed allylation of imine **5a**.^[a]



Entry	[Rh] (2.5 mol%)	Ligand (5 mol%)	Conv [%] ^[b]	ee [%] ^[c]
1	$[\text{Rh}(\text{cod})\text{Cl}]_2$	none	>95 ^[d]	–
2	$[\text{Rh}(\text{C}_2\text{H}_4)_2\text{Cl}]_2$	L1	<5	–
3	$[\text{Rh}(\text{C}_2\text{H}_4)_2\text{Cl}]_2$	L2	60	67
4	$[\text{Rh}(\text{L3})\text{Cl}]_2$	none	>95	93

[a] Reactions were conducted using 0.10 mmol of **5a**. [b] Determined by ¹H NMR analysis of the unpurified reaction mixtures. [c] Determined by HPLC analysis on a chiral stationary phase. [d] *rac*-**6a** isolated in 87% yield.

[*] Dr. Y. Luo, H. B. Hepburn, N. Chotsaeng, Dr. H. W. Lam
EaStCHEM, School of Chemistry, University of Edinburgh
Joseph Black Building, The King's Buildings
West Mains Road, Edinburgh EH9 3JJ, United Kingdom
E-mail: h.lam@ed.ac.uk
Homepage: <http://homepages.ed.ac.uk/hlam/index.html>

[**] We thank the ERC (Starting Grant No. 258580), the EPSRC (Leadership Fellowship to H. W. L.), and the Thai government for support of this work, Dr. Gary S. Nichol (University of Edinburgh) for X-ray crystallography, and the EPSRC National Mass Spectrometry Service Centre at the University of Wales, Swansea, for high-resolution mass spectra. We thank Dr. Uwe Schneider (University of Edinburgh) for helpful discussions.



consumption of **5a** was observed after 3 h to give **6a** in 87% isolated yield (entry 1). Next, the use of chiral ligands was examined. While (*R*)-BINAP (**L1**) was totally ineffective in promoting the reaction (entry 2), chiral diene **L2**^[17, 18] provided (*R*)-**6a** in 60% conversion with a promising 67% *ee* (entry 3). However, using 2.5 mol% of the rhodium complex derived from chiral diene **L3**,^[16, 19] (*R*)-**6a** was obtained in >95% conversion and 93% *ee* (entry 4).^[20] The results of Tables 1 and 2 clearly highlight the benefits of a cyclic imine structure in facilitating efficient allylation.^[21] It should be noted that the use of potassium allyltrifluoroborate (**2a**) was essential for high enantioselectivity; repeating the reaction in Table 2, entry 4 using allylboronic acid pinacol ester in place of **2a**, with the addition of aqueous K₃PO₄ (0.5 equiv), led to

Table 3: Enantioselective Rh-catalyzed allylation of cyclic imines.

Entry	Product		Yield [%] ^[a]	<i>ee</i> [%] ^[b]
1		6a R = H	87	96
2		6b R = Me	93	97
3		6c R = OMe	92	91
4 ^[c]		6d R = F	86	98
5 ^[c]		6e R = Cl	96	96
6 ^[c]		6f R = Br	79	94
7 ^[c]		6g R = CN	97	98
8		6h	>95	95
9		6i	94	98
10		6j	95	93
11 ^[d]		6k R = H	59	95
12 ^[d]		6l R = Cl	65	94
13 ^[d]		6m R = Br	45	90
14		6n R = Me	84	99 ^[e]
15 ^[f]		6o R = Ph	88	97
16 ^[g]		6p	83	93 ^[e]

[a] Isolated yields. [b] Determined by HPLC analysis on a chiral stationary phase. [c] Using *i*PrOH (13 equiv) in toluene/dioxane for a reaction time of 18 h. [d] Reaction time of 15 h. [e] Enantiomeric excess determined on the *N*-benzyl derivative. [f] Reaction time of 12 h. [g] Using dioxane in place of THF, at 80 °C, for 15 h.

(*R*)-**6a** in >95% conversion, but in only 28% *ee*. The reasons for this lower selectivity are not clear at the present time.

With an effective ligand identified, the scope of this process was investigated under slightly modified conditions (Table 3). Using 1.5 mol% of [$\text{Rh}(\text{L3})\text{Cl}$]₂ and MeOH (5 equiv) in a THF/dioxane mixture at 55 °C, a wide range of cyclic imines underwent highly enantioselective allylations (90–99% *ee*) in generally good yields. Benzoxathiazine-2,2-dioxides containing various substituents (methyl, methoxy, halogen, cyano, or dioxole) were effective substrates (entries 1–10), though in certain cases, the use of *i*PrOH (13 equiv) in toluene/dioxane rather than MeOH in THF/dioxane was required to maintain high enantioselectivities (entries 4–7). Other cyclic aldimines such as 1,2,6-thiadiazine-1,1-dioxides also underwent allylation (entries 11–13), though yields were lower in these cases. The process is not restricted to aldimines; cyclic ketimines such as 1,2,6-thiadiazolidine 1,1-dioxides (entries 14 and 15) and a cyclic sulfamidate imine (entry 16) are also viable substrates. These results (entries

Table 4: Allylation of cyclic imines with various allyltrifluoroborates.^[a]


Allyltrifluoroborate	Product using 5a	Product using 5o
	 7a 91% 15:1 d.r., 99% <i>ee</i>	 7e 68% >19:1 d.r., 97% <i>ee</i>
	 7b 89% ^[b] 6:1 d.r., 99% <i>ee</i>	 7f 89% >19:1 d.r., 99% <i>ee</i> ^[c]
	 7c 76% ^[b] 11:1 d.r., 99% <i>ee</i>	 7g 61% ^[b] 17:1 d.r., 99% <i>ee</i>
	 7d 91% 97% <i>ee</i>	 7h 75% 95% <i>ee</i>

[a] Cited yields are of isolated pure major diastereomers (where relevant). Diastereomeric ratios were determined by ¹H NMR analysis of the unpurified reaction mixtures. Enantiomeric excesses were determined by HPLC analysis on a chiral stationary phase. [b] Isolated yield of an inseparable mixture of diastereomers. [c] Reaction time of 12 h.

14–16) are notable because catalytic enantioselective allylations of ketimines are rare.^[9g,12a,b]


Table 4 presents the allylation of cyclic imines **5a** and **5o**^[22] with more highly substituted potassium allyltrifluoroborates **2b–e**. Here, clean allylic transposition took place to form a new carbon–carbon bond at the more substituted γ -carbon of the allyltrifluoroborate. *E*-Crotyltrifluoroborate **2b** underwent highly enantioselective additions to **5a** and **5o** to afford *anti* products **7a** and **7e**, respectively, with high diastereoselectivities. Importantly, *Z*-crotyltrifluoroborate **2c** afforded the corresponding *syn* products **7b** and **7f** with high enantioselectivities, though the diastereomeric ratio in the case of **7b** was more modest (6:1 d.r.). Similarly to **2b**, *E*-2-hexen-1-yltrifluoroborate **2d** afforded *anti* products with high levels of diastereo- and enantioselection (products **7c** and **7g**). Finally, prenyltrifluoroborate **2e** was also highly effective, reacting with aldimine **5a** to deliver reverse prenylation product **7d** from aldimine **5a** in 97% *ee*, and with ketimine **5o** to give **7h** containing two vicinal quaternary centers in 95% *ee*.

To gain some insight into the mechanism of these reactions, allylation of ketimine **5o** with dideuterated potassium allyltrifluoroborate **8** was conducted [Eq. (1)].^[23] This reaction resulted in a 1:1 mixture of products **9a** and **9b** in 88% yield, which suggests that allylation proceeds via the intermediacy of an allylrhodium(I) species that undergoes rapid interconversion between the two σ -allyl haptomers to scramble the deuterium label.^[24] Alternative mechanisms involving an allylboron species as the reactive intermediate would be expected to furnish **9b** only.^[8]



Since *E*- and *Z*-crotyltrifluoroborates **2b** and **2c** provided *anti* and *syn* allylation products, respectively, with good to high diastereoselectivities (Table 2), it is likely that: (i) cyclic 6-membered transition states are operative, and (ii) the corresponding allylrhodium(I) intermediates possess appreciable configurational stability during the time scale of allylation despite the possibility of rapid interconversion between the two σ -allyl isomers which could erode *E/Z* stereochemistry.^[24,25]

On the basis of these features, a possible catalytic cycle for these reactions, using imine **5a** for illustrative purposes, is shown in Scheme 1. Presumably, in the presence of MeOH, potassium allyltrifluoroborates can undergo reversible methanolysis to generate a mixed alkoxide/fluoride boron ate-complex **10**,^[26] which engages in transmetalation with the chiral diene-ligated rhodium complex **11** (where X = Cl, F, or OMe) to form the allylrhodium species **12**. Coordination of



Scheme 1. Possible catalytic cycle.

the imine **5a** then occurs as shown in **13** to minimize unfavorable steric interactions between the imine activating group and a phenyl substituent of the ligand (as would occur in the disfavored diastereomeric complex **14**). Allylation of the bound imine in **13** through a cyclic chairlike transition state would give **15**, which is then protonated with HX (X = Cl, F, or OMe) to release the product and regenerate **11**.

In conclusion, a chiral diene-ligated rhodium complex has been shown to be highly effective in catalyzing the addition of potassium allyltrifluoroborates to cyclic aldimines and ketimines. To our knowledge, these reactions represent the first rhodium-catalyzed enantioselective nucleophilic allylations of π -electrophiles with allylboron compounds, and proceed with generally good yields and high levels of diastereo- and enantioselection. Where relevant, the chiral allylrhodium(I) intermediates possess appreciable *E/Z* configurational stability during the time scale of allylation, allowing access to different product diastereomers through the use of *E*- or *Z*-allyltrifluoroborates. Further investigations of enantioselective rhodium-catalyzed nucleophilic allylations are underway, and the results of these studies will be reported in due course.

Received: ((will be filled in by the editorial staff))
Published online on ((will be filled in by the editorial staff))

Keywords: allyltrifluoroborate • asymmetric catalysis • enantioselectivity • imines • rhodium


- [1] For relevant reviews, see: a) P. Tian, H.-Q. Dong, G.-Q. Lin, *ACS Catalysis* **2012**, 2, 95–119; b) H. J. Edwards, J. D. Hargrave, S. D. Penrose, C. G. Frost, *Chem. Soc. Rev.* **2010**, 39, 2093–2105; c) C. Defieber, H. Grutzmacher, E. M. Carreira, *Angew. Chem.* **2008**, 120, 4558–4579; *Angew. Chem. Int. Ed.* **2008**, 47, 4482–4502; d) J. B. Johnson, T. Rovis, *Angew. Chem.* **2008**, 120, 852–884; *Angew. Chem. Int. Ed.* **2008**, 47, 840–871; e) K. Yoshida, T. Hayashi, *Modern Rhodium-Catalyzed Organic Reactions* **2005**, 55–77; f) T. Hayashi, K. Yamasaki, *Chem. Rev.* **2003**, 103, 2829–2844.

- [2] For representative examples, see: a) Y. Takaya, M. Ogasawara, T. Hayashi, M. Sakai, N. Miyaara, *J. Am. Chem. Soc.* **1998**, *120*, 5579-5580; b) Y. Takaya, M. Ogasawara, T. Hayashi, *Tetrahedron Lett.* **1998**, *39*, 8479-8482; c) T. Hayashi, K. Ueyama, N. Tokunaga, K. Yoshida, *J. Am. Chem. Soc.* **2003**, *125*, 11508-11509; d) N. Tokunaga, T. Hayashi, *Adv. Synth. Catal.* **2007**, *349*, 513-516; e) J. L. Zigterman, J. C. S. Woo, S. D. Walker, J. S. Tedrow, C. J. Borths, E. E. Bunel, M. M. Faul, *J. Org. Chem.* **2007**, *72*, 8870-8876; e) R. Shintani, Y. Ichikawa, K. Takatsu, F.-X. Chen, T. Hayashi, *J. Org. Chem.* **2008**, *74*, 869-873; f) V. Hickmann, A. Kondoh, B. Gabor, M. Alcarazo, A. Fürstner, *J. Am. Chem. Soc.* **2011**, *133*, 13471-13480; g) T. Thaler, L.-N. Guo, A. K. Steib, M. Raducan, K. Karaghiosoff, P. Mayer, P. Knochel, *Org. Lett.* **2011**, *13*, 3182-3185.
- [3] S. Crotti, F. Bertolini, F. Macchia, M. Pineschi, *Chem. Commun.* **2008**, 3127-3129.
- [4] For rhodium-catalyzed asymmetric alkynylation using terminal alkynes, see: a) P. K. Dhondt, P. Carberry, L. B. Choi, J. D. Chisholm, *J. Org. Chem.* **2007**, *72*, 9590-9596; b) T. Nishimura, X.-X. Guo, N. Uchiyama, T. Katoh, T. Hayashi, *J. Am. Chem. Soc.* **2008**, *130*, 1576-1577; c) T. Nishimura, E. Tsurumaki, T. Kawamoto, X.-X. Guo, T. Hayashi, *Org. Lett.* **2008**, *10*, 4057-4060; d) E. Fillion, A. K. Zoritto, *J. Am. Chem. Soc.* **2009**, *131*, 14608-14609; e) T. Nishimura, T. Sawano, S. Tokui, T. Hayashi, *Chem. Commun.* **2010**, 46, 6837-6839; f) T. Ohshima, T. Kawabata, Y. Takeuchi, T. Kakinuma, T. Iwasaki, T. Yonezawa, H. Murakami, H. Nishiyama, K. Mashima, *Angew. Chem.* **2011**, *123*, 6420-6424; *Angew. Chem. Int. Ed.* **2011**, *50*, 6296-6300.
- [5] For enantioselective rhodium-catalyzed alkynylation of enones using alkynylsilanols, see: T. Nishimura, S. Tokui, T. Sawano, T. Hayashi, *Org. Lett.* **2009**, *11*, 3222-3225.
- [6] For rhodium-catalyzed isomerization of alkenylboron reagents into allylboron reagents followed by *in situ* racemic allylation of aldehydes, see: H. Shimizu, T. Igarashi, T. Miura, M. Murakami, *Angew. Chem.* **2011**, *123*, 11667-11671; *Angew. Chem. Int. Ed.* **2011**, *50*, 11465-11469.
- [7] For rhodium-catalyzed redox allylation of ketones with allyl acetate and bis(pinacolato)diboron, see: F. J. Williams, R. E. Grote, E. R. Jarvo, *Chem. Commun.* **2012**, 48, 1496-1498.
- [8] a) S. E. Denmark, J. P. Fu, *Chem. Rev.* **2003**, *103*, 2763-2793; b) M. Yus, J. González-Gómez, F. Foubelo, *Chem. Rev.* **2011**, *111*, 7774-7854.
- [9] To our knowledge, enantioselective rhodium-catalyzed nucleophilic allylations are limited to additions of allylstannanes to aldehydes (where a chiral Rh(III) complex functions as a Lewis acid), and cyclizations of allylrhodium species generated by additions to allenes. See: a) Y. Motoyama, H. Narusawa, H. Nishiyama, *Chem. Commun.* **1999**, 131-132; b) M. Shi, G. X. Lei, Y. Masaki, *Tetrahedron: Asymmetry* **1999**, *10*, 2071-2074; c) Y. Motoyama, M. Okano, H. Narusawa, N. Makihara, K. Aoki, H. Nishiyama, *Organometallics* **2001**, *20*, 1580-1591; d) Y. Motoyama, H. Nishiyama, *Synlett* **2003**, 1883-1885; e) Y. Motoyama, T. Sakakura, T. Takemoto, K. Shimozono, K. Aoki, H. Nishiyama, *Molecules* **2011**, *16*, 5387-5401; f) X.-X. Guo, T. Sawano, T. Nishimura, T. Hayashi, *Tetrahedron: Asymmetry* **2010**, *21*, 1730-1736; g) D. N. Tran, N. Cramer, *Angew. Chem.* **2010**, *122*, 8357-8360; *Angew. Chem. Int. Ed.* **2010**, *49*, 8181-8184.
- [10] T. R. Ramadhar, R. A. Batey, *Synthesis* **2011**, 1321-1346.
- [11] For representative examples of catalytic enantioselective allylation of imines or their derivatives with allyl nucleophiles other than allylboron reagents, see: a) H. Nakamura, K. Nakamura, Y. Yamamoto, *J. Am. Chem. Soc.* **1998**, *120*, 4242-4243; b) T. Gastner, H. Ishitani, R. Akiyama, S. Kobayashi, *Angew. Chem.* **2001**, *113*, 1949-1951; *Angew. Chem. Int. Ed.* **2001**, *40*, 1896-1898; c) S. Kobayashi, C. Ogawa, H. Konishi, M. Sugiura, *J. Am. Chem. Soc.* **2003**, *125*, 6610-6611; d) R. A. Fernandes, A. Stimac, Y. Yamamoto, *J. Am. Chem. Soc.* **2003**, *125*, 14133-14139; e) R. A. Fernandes, Y. Yamamoto, *J. Org. Chem.* **2004**, *69*, 735-738; f) C. Ogawa, M. Sugiura, S. Kobayashi, *Angew. Chem.* **2004**, *116*, 6653-6655; *Angew. Chem. Int. Ed.* **2004**, *43*, 6491-6493; g) H. Kiyohara, Y. Nakamura, R. Matsubara, S. Kobayashi, *Angew. Chem.* **2006**, *118*, 1645-1647; *Angew. Chem. Int. Ed.* **2006**, *45*, 1615-1617; h) R. Kargbo, Y. Takahashi, S. Bhor, G. R. Cook, G. C. Lloyd-Jones, I. R. Shepperson, *J. Am. Chem. Soc.* **2007**, *129*, 3846-3847; i) K. L. Tan, E. N. Jacobsen, *Angew. Chem.* **2007**, *119*, 1337-1339; *Angew. Chem. Int. Ed.* **2007**, *46*, 1315-1317; j) X.-C. Qiao, S.-F. Zhu, W.-Q. Chen, Q.-L. Zhou, *Tetrahedron: Asymmetry* **2010**, *21*, 1216-1220.
- [12] a) R. Wada, T. Shibuguchi, S. Makino, K. Oisaki, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* **2006**, *128*, 7687-7691; b) M. Kanai, R. Wada, T. Shibuguchi, M. Shibasaki, *Pure Appl. Chem.* **2008**, *80*, 1055-1062; c) E. M. Vieira, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* **2011**, *133*, 3332-3335; d) J. Aydin, K. S. Kumar, M. J. Sayah, O. A. Wallner, K. Szabó, *J. Org. Chem.* **2007**, *72*, 4689-4697; e) M. Fujita, T. Nagano, U. Schneider, T. Hamada, C. Ogawa, S. Kobayashi, *J. Am. Chem. Soc.* **2008**, *130*, 2914-2915; f) A. Chakrabarti, H. Konishi, M. Yamaguchi, U. Schneider, S. Kobayashi, *Angew. Chem.* **2010**, *122*, 1882-1885; *Angew. Chem. Int. Ed.* **2010**, *49*, 1838-1841; g) Y.-Y. Huang, A. Chakrabarti, N. Morita, U. Schneider, S. Kobayashi, *Angew. Chem.* **2011**, *123*, 11317-11320; *Angew. Chem. Int. Ed.* **2011**, *50*, 11121-11124; h) S. Lou, P. N. Moquist, S. E. Schaus, *J. Am. Chem. Soc.* **2007**, *129*, 15398-15404.
- [13] For the catalytic enantioselective diboration of allenes, followed by diastereoselective additions of the resulting allylboronic esters to imines, see: J. D. Sieber, J. P. Morken, *J. Am. Chem. Soc.* **2005**, *128*, 74-75.
- [14] For the first transition-metal-catalyzed additions of potassium allyltrifluoroborate to imines, see: N. Solin, O. A. Wallner, K. J. Szabó, *Org. Lett.* **2005**, *7*, 689-691.
- [15] For reviews on organotrifluoroborates, see: a) G. A. Molander, D. L. Sandrock, *Curr. Opin. Drug Discovery Dev.* **2009**, *12*, 811-823; b) S. Darses, J.-P. Genet, *Chem. Rev.* **2008**, *108*, 288-325; c) H. A. Stefani, R. Cella, A. S. Vieira, *Tetrahedron* **2007**, *63*, 3623-3658; d) G. A. Molander, R. Figueroa, *Aldrichim. Acta* **2005**, *38*, 49-56.
- [16] Y. Luo, A. J. Carnell, H. W. Lam, *Angew. Chem.* **2012**, *124*, Early View. DOI: 10.1002/ange.201202136; *Angew. Chem. Int. Ed.* **2012**, *51*, Early View. DOI: 10.1002/anie.201202136.
- [17] K. Okamoto, T. Hayashi, V. H. Rawal, *Chem. Commun.* **2009**, 4815-4817.
- [18] For reviews of chiral diene ligands, see: a) C.-G. Feng, M.-H. Xu, G.-Q. Lin, *Synlett* **2011**, 1345-1356; b) R. Shintani, T. Hayashi, *Aldrichim. Acta* **2009**, *42*, 31-38; c) J. B. Johnson, T. Rovis, *Angew. Chem.* **2008**, *120*, 852-884; *Angew. Chem. Int. Ed.* **2008**, *47*, 840-871; d) C. Defieber, H. Grutzmacher, E. M. Carreira, *Angew. Chem.* **2008**, *120*, 4558-4579; *Angew. Chem. Int. Ed.* **2008**, *47*, 4482-4502.
- [19] Y. Luo, A. J. Carnell, *Angew. Chem.* **2010**, *122*, 2810-2814; *Angew. Chem. Int. Ed.* **2010**, *49*, 2750-2754.
- [20] The relative and absolute configurations of the products obtained herein were assigned by analogy with those of **6h**, **6i**, and **7d** which were determined by X-ray crystallography. See Supporting Information for details. CCDC 880900-880902. contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [21] For application of a catalytic enantioselective addition of an allylsilane to a cyclic imine in total synthesis, see: T. Itoh, M. Miyazaki, H. Fukuoka, K. Nagata, A. Ohsawa, *Org. Lett.* **2006**, *8*, 1295-1297.
- [22] Attempted allylation of cyclic imines **5k-m** with more substituted allyltrifluoroborate reagents provided complex mixtures of products, while imine **5p** was unreactive toward these reagents.
- [23] For similar experiments, see: a) J. D. Sieber, J. P. Morken, *J. Am. Chem. Soc.* **2008**, *130*, 4978-4983; b) T. J. Barker, E. R. Jarvo, *Org. Lett.* **2009**, *11*, 1047-1049; c) U. Schneider, H. T. Dao, S. Kobayashi, *Org. Lett.* **2010**, *12*, 2488-2491.
- [24] For a discussion of the enyl ($\sigma + \pi$) character of electrophilic allylrhodium(III) species, see: P. A. Evans, J. D. Nelson, *J. Am. Chem. Soc.* **1998**, *120*, 5581-5582.
- [25] Trost, B. M.; VanVranken, D. L. *Chem. Rev.* **1996**, *96*, 395-422.
- [26] For a discussion of organotrifluoroborate hydrolysis in Suzuki-Miyaura reactions, see: a) M. Butters, J. N. Harvey, J. Jover, A. J. J. Lennox, G. C. Lloyd-Jones, P. M. Murray, *Angew. Chem.* **2010**, *122*, 5282-5286; *Angew. Chem. Int. Ed.* **2010**, *49*, 5156-5160; b) A. J. J. Lennox, G. C. Lloyd-Jones, *J. Am. Chem. Soc.* **2012**, *134*, 7431-7441.

Asymmetric Allylation

Y. Luo, H. B. Hepburn, N. Chotsaeng,
H. W. Lam* **Page – Page**

Enantioselective Nucleophilic Allylation
of Cyclic Imines with Allylboron
Reagents Using a Chiral Diene-Ligated
Rhodium Catalyst



Chiral allylrhodium nucleophiles: In the presence of a chiral diene-ligated rhodium complex, a range of cyclic imines undergo highly diastereo- and enantioselective allylations with various potassium allyltrifluoroborates. These

reactions most likely proceed via allylrhodium(I) intermediates, and represent the first rhodium-catalyzed enantioselective nucleophilic allylations of π -electrophiles with allylboron compounds.